



EXCISION OF MUCOCELE WITH GOLDMAN-FOX SOFT TISSUE NIPPER

EXCISÃO DE MUCOCELE COM TESOURA DE TECIDOS GOLDMAN-FOX

ESCISIÓN DE MUCOCELO CON PINZA DE TEJIDO BLANDO GOLDMAN-FOX

Diego Leonardo de Souza¹, Edcléia Leite Brandão dos Santos², Gustavo Souza Alves³, Vitor Rizzotti Galvão⁴, Cleomar Donizeth Rodrigues⁵, Victor Milis Wandelli⁶, Virgílio Pimentel Galvão⁷

e432856

<https://doi.org/10.47820/recima21.v4i3.2856>

RECEIVED: 01/14/2023

APPROVED: 02/14/2023

PUBLISHED: 03/05/2023

ABSTRACT

Purpose: This report aims to describe a technique for surgical removal of lip mucocèles with Goldman-Fox soft tissue nipper and analysis the clinical aspects. **Materials and Methods:** Mucocèle of the lower lip was completely removed with a Goldman-Fox soft tissue nipper. The lesion was sent to histopathological analysis, which confirmed the clinical diagnosis of mucocèle. **Results:** After excision, there was very minimal *bleeding and* regular margins were obtained, which contributed for suture and soft repair. The lesion was completely removed with no extravasations of mucous. There was no recurrence of the lesion after 2-years follow-up. **Conclusions:** The excision with Goldman-Fox soft tissue nipper to treat the mucocèle of the lower lip was effective and had not complications, with shorter operative time, which is especially good for children and for less cooperative patients. Besides, this technique allowed the histopathological analysis to confirm the diagnosis.

KEYWORDS: Mucocèle. Excision. Lower lip. Goldman-Fox Soft Tissue Nipper.

RESUMO

Objetivo: Este relato teve como objetivo descrever uma técnica para remoção cirúrgica de mucocèle labial com pinça de Goldman-Fox para tecidos moles e analisar os aspectos clínicos. **Materiais e Métodos:** A mucocèle do lábio inferior foi totalmente removida com pinça de tecido mole Goldman-Fox. A lesão foi encaminhada para análise histopatológica, que confirmou o diagnóstico clínico de mucocèle. **Resultados:** Após a excisão, houve sangramento mínimo e margens regulares foram obtidas, o que contribuiu para sutura e reparação suave. A lesão foi totalmente removida sem extravasamento de mucosa. Não houve recidiva da lesão após 2 anos de preservação. **Conclusões:** A excisão com pinça de Goldman-Fox para tratar a mucocèle do lábio inferior foi eficaz e sem complicações, com menor tempo operatório, o que é especialmente bom para crianças e para pacientes menos cooperativos. Além disso, esta técnica permitiu a análise histopatológica para confirmar o diagnóstico.

PALAVRAS-CHAVE: Mucocèle. Excisão. Lábio inferior. Pinça de tecidos moles Goldman-Fox.

RESUMEN

Objetivo: Este informe tiene como objetivo describir una técnica para la extracción quirúrgica de mucocèles labiales con fórceps de Goldman-Fox para tejidos blandos y analizar los aspectos clínicos. **Materiales y Métodos:** El mucocèle del labio inferior se eliminó por completo con pinzas para tejidos blandos Goldman-Fox. Se envió la lesión para análisis histopatológico, que confirmó el diagnóstico clínico de mucocèle. **Resultados:** Después de la extirpación, hubo sangrado mínimo y se obtuvieron márgenes regulares, lo que contribuyó a la sutura y reparación suave. La lesión se eliminó por completo sin extravasación de la mucosa. No hubo recurrencia de la lesión después de 2

¹ Instituto Orofacial das Américas.

² FACIPLAC.

³ Odontodoc.

⁴ Instituto Aria.

⁵ UNICEPLAC – Centro Universitário do Planalto Central Aparecido dos Santos.

⁶ Cirurgião-dentista (Universidade Federal de Santa Catarina).

⁷ Faculdade Odontologia da USP – FOUASP.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EXCISION OF MUCOCELE WITH GOLDMAN-FOX SOFT TISSUE NIPPER
Diego Leonardo de Souza, Edcléia Leite Brandão dos Santos, Gustavo Souza Alves, Vitor Rizzotti Galvão,
Cleomar Donizeth Rodrigues, Victor Milis Wandelli, Virgílio Pimentel Galvão

años de seguimiento. Conclusiones: La escisión con fórceps de Goldman-Fox para el tratamiento del mucocele del labio inferior fue eficaz y sin complicaciones, con un tiempo operatorio más corto, lo que es especialmente bueno para los niños y para los pacientes menos cooperativos. Además, esta técnica permitió el análisis histopatológico para confirmar el diagnóstico.

PALABRAS CLAVE: Mucocele. Excisión. Labio inferior. Pinzas para tejidos blandos de Goldman-Fox.

INTRODUCTION

Mucoceles are common benign lesions of the oral mucosa which mechanisms for mucous cavity development are extravasation or retention¹. Extravasation is the leakage of fluid from the salivary ducts into the surrounding tissue and is the primary cause of mucocele formation with physical trauma as the initiating factor¹. Retention phenomenon occurs as a result of a narrowed ductal opening that cannot adequately accommodate the exit of saliva produced, leading to ductal dilation and surface swelling which results from rupture of a salivary gland duct¹. The lesion mainly affects young people, and no gender difference was observed¹⁻³. Regarding the location of the mucoceles, the lower lip is the most prevalent²⁻⁴ because it is the area most often susceptible to trauma⁵. Improper treatment can result in a recurrence of the lesion⁴.

Several treatments have been proposed in the literature, such as excision of the lesion associated or not with removal of the gland involved, intralesional injection therapy with OK-432, carbon dioxide laser, cryosurgery and micro-marsupialization⁵⁻¹⁰. Non surgical techniques for the treatment of mucocele has been widely used and reported as effective techniques, with less discomfort and recommended primarily for non-cooperative patients^{5,6,8}. In these cases where there is no surgical removal of the lesion, the histopathological examination may not be executed and, therefore, the treatment is performed without confirmation of definitive diagnosis. Histopathological analysis is critical, as some malignant lesions are clinically similar to oral mucoceles¹¹⁻¹³.

The aim of this report was to describe a technique for surgical removal of lip mucoceles with Goldman-Fox soft tissue nipper, describing the clinical and histopathological aspects.

CASE REPORT

A 14-year-old female patient was referred with lower lip lesion associated with periodically swelling during the last three months. The patient has a history of biting the lip frequently. Clinical examination revealed the presence of an exofitic lesion in the lower lip with 6mm of diameter, sessile, soft consistency, asymptomatic and color of adjacent normal mucosa (Figure 1). All the other structures were normal at the oral clinical exam.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EXCISION OF MUCOCELE WITH GOLDMAN-FOX SOFT TISSUE NIPPER
Diego Leonardo de Souza, Edcléia Leite Brandão dos Santos, Gustavo Souza Alves, Vitor Rizzotti Galvão,
Cleomar Donizeth Rodrigues, Victor Milis Wandelli, Virgílio Pimentel Galvão



Figure 1. Exofitic lesion on the lower lip mucosa with approximately 6 mm of diameter.

SURGICAL TECHNIQUE AND POSTOPERATIVE CARE

Aqueous solution of Chlorhexidine 0,12% was used to disinfect and perform surgical antisepsis. Local anesthesia was achieved by mental nerve block using 2% Lidocain with epinephrine (1:100.000). The Goldman-Fox soft tissue nipper (Hu-Friedy) was open positioned around the lesion on its basis (Figure 2). It was necessary just a movement for excision (Figure 3) and the lesion was completely removed without extravasations of mucus (Figure 4).



Figure 2. Placement of Goldman-Fox Soft tissue nipper on the basis of mucosa around the lesion.



Figure 3. Move with single cut.



Figure 4. Excision performed on the basis of lesion with the closure of the Goldman-Fox soft tissue nipper.

Light bleeding from the surgical wound occurred after the excision and regular margins were observed (Figure 5). This characteristic permits the proper alignment of the *wound* edges and *contributes* to their final suture. The lesion was fixed in *10% buffered formalin*.



Figure 5. Suture performed.



Figure 6. The lesion was removed entirely: edges of the ellipse-shaped surgical wound, which occurred after use of cutting instrument.

The patient was examined at 1 week after surgery and there were no complaints of swelling, bleeding or pain.

DISCUSSION

Mucocele is a benign lesion of salivary gland, and the primary treatment modality consists basically in surgical removal and can be performed with various methods⁷⁻¹⁰. In the literature, different treatments are reported according to the size of the lesion⁵⁻¹⁰ and small, superficial lesions may have spontaneous resolution without necessity of treatment^{3, 14}. Cryosurgery is an alternative method of treatment which aims for minimal injury to adjacent structures¹⁵ but is associated with a significant number of recurrences that require reoperation with classic or alternative surgical procedures⁸ and the technique alone do not permit histopathological analysis.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EXCISION OF MUCOCELE WITH GOLDMAN-FOX SOFT TISSUE NIPPER
Diego Leonardo de Souza, Edcléia Leite Brandão dos Santos, Gustavo Souza Alves, Vitor Rizzotti Galvão,
Cleomar Donizeth Rodrigues, Victor Milis Wandelli, Virgílio Pimentel Galvão

The treatment with soft tissue laser offers an alternative treatment with simple mucocele ablation and short surgical time. The wound remains without suture and a little discomfort was reported^{6, 16}. The laser does not remove but only vaporizes the lesion and perilesional tissue which does not make possible the histopathological analysis of the lesion after the procedure. Some authors have reported an incisional biopsy before the use of carbon dioxide laser⁶, therefore the conventional technique with bistoury cannot be waived. Besides, the equipment is expensive and adequate measures must be taken for both the patient and the staff's protection.⁶

In relation to micro-dissection technique, microsurgical instruments contribute the surgeon to dissect the mucocele¹, however this procedure spends relative high time. Care should be taken to avoid injury to the other glands and ducts with the suture needle because surgical injury to any of the marginal glands is another cause of recurrence¹.

The recently introduced intralesional injection therapy with OK-432 was reported,⁵ however it seems to occur high recurrence of mucocele. Besides, the technique does not permit the histopathological analysis. This is an important point to consider since adenocarcinomas mimicking clinical aspects of mucocele have been reported^{11-13, 17, 18}.

In our case report, the used of surgical excision with Goldman-Fox soft tissue nipper was used to treat a mucocele of 6mm. This technique is recommended to treat primarily small to medium-sized lesions. Large mucoceles (size > 10mm) are best treated with an unroofing procedure (marsupialization), because excision or dissection would be problematic and risk vital structures such as the labial branch of the mental nerve¹. A clinicopathologic review of 1,824 cases of mucoceles reported that the maximum diameter at mean was 8mm⁴; therefore, the Goldman-Fox soft tissue nipper may be indicated for lesions until this size because the cutting blade diameter has 8mm. For excision of a minor injury, simply position the pliers so closer to its end, delimiting it for removal. The advantages include a bloodless operating field, precise incision, easy surgery, decreased postoperative swelling, low cost and minimized scar tissue. No particular complications in the postoperative period and no hemorrhagic episodes occurred.

It has been suggested to observe the lesion for 5 months before proceeding with surgical treatment because this is an adequate time for spontaneous resolution of the lesion¹⁰. A 2 year follow-up was done and no recurrence was verified.

CONCLUSION

This is the first case reporting the excision with Goldman-Fox soft tissue nipper to treat mucocele of the lower lip which was effective and had not complications. The operative time is shorter than with the conventional excisional methods, which is especially good for children and for less cooperative patients. Besides, this technique allowed the histopathological analysis to confirm the diagnosis.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR

ISSN 2675-6218

EXCISION OF MUCOCELE WITH GOLDMAN-FOX SOFT TISSUE NIPPER
 Diego Leonardo de Souza, Edcléia Leite Brandão dos Santos, Gustavo Souza Alves, Vitor Rizzotti Galvão,
 Cleomar Donizeth Rodrigues, Victor Milis Wandelli, Virgílio Pimentel Galvão

REFERENCES

1. Baumash HD. Mucoceles and Ranulas. *J Oral Maxillofac Surg.* 2003;61:369-378.
2. Re Cecconi D, Achilli A, Tarozzi M, Lodi G, Demarosi F, Sardella A, et al. Mucoceles of the oral cavity: a large case series (1994-2008) and a literature review. *Med Oral Patol Oral Cir Bucal.* 2010;15:e551-6.
3. Minguéz IM, Bonet CC, Ata-Ali JM, Carrillo CG, MD, Peñarrocha MD. Clinical characteristics, treatment, and evolution of 89 mucoceles in children. *J Oral Maxillofac Surg.* 2010;68:2468-2471.
4. Chi AC, Lambert PR 3rd, Richardson MS, Neville BW. Oral mucoceles: a clinicopathologic review of 1,824 cases, including unusual variants. *J Oral Maxillofac Surg.* 2011;69:1086-93.
5. Ohta N, Fukase S, Suzuki Y, Aoyagi M. Treatment of salivary mucocele of the lower lip by OK-432. *Auris Nasus Larynx.* 2011;38:240-243
6. Huang IY, Chen CM, Kao YH, Worthington P. Treatment of Mucocele with Carbon Dioxide Laser. *J Oral Maxillofac Surg.* 2007;65:855-858.
7. Tilaveridis I, Lazaridou M, Lazaridis N. Excision of Lower Lip Mucocele. *J Oral Maxillofac Surg.* 2011;69:1408-1410.
8. Marcushamer M, King DL, Ruano NS. Cryosurgery in the management of mucoceles in children. *Pediatr Dent.* 1997 May-Jun;19(4):292-3.
9. Delbem AC, Cunha RF, Vieira AE, Ribeiro LL. Treatment of mucus retention phenomena in children by the micro-marsupialization technique: case reports. *Pediatr Dent.* Mar-2000 Apr;22(2):155-8.
10. Pandit RT, Park AH. Management of pediatric ranula. *Otolaryngol Head Neck Surg.* 2002;127:115-8.
11. Omlie JE, Koutlas IG. Acinic Cell Carcinoma of Minor Salivary Glands: a clinicopathologic study of 21 cases. *J Oral Maxillofac Surg.* 2010;68:2053-2057.
12. Madhavan NR, Ramchandran CR, Veeravama V. Canalicular adenoma of the upper lip mimicking mucocele. *Indian J Dent Res.* 2004;15:66-69.
13. Closmann JJ, Torske KR. When a mucocele is not mucocele: adenocarcinoma NOS- a case report and review of the literature. *Gen Dent.* 2007;55:325-327.
14. Bodner L, Manor E, Joshua BZ, Shaco-Levy R. Oral Mucoceles in Children- Analysis of 56 New Cases. *Pediatr Dermatol.* 2015 Sep-Oct;32(5):647-50.
15. Aulak h, et al. Cryotherapy for treatment of mucocele. *Niger J Surg.* 2016 Jul-Dec; 22(2):130-133.
16. Ramkumar S, Ramkumar L, Malathi N, Suganya R. Excision of Mucocele Using Diode Laser in Lower Lip. *Case Rep Dent.* 2016:1746316.
17. Tandon A, Sircar K, Chowdhry A, Bablani D. Salivary duct cyst on lower lip: A rare entity and literature review. *J Oral Maxillofac Pathol.* 2014 Sep;18(1):151-156.
18. Karthikeyan M, Varghese A, Vasupradha G, Dinakaran J. Mucocele: A diagnostic dilemma!! *J Pharm Bioallied Sci.* 2016 Oct;8(1):168-170.