



DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL

DIVERSIDADE DE FORMIGAS (HYMENOPTERA: FORMICIDAE) OCORRENTES PARA O AMAPÁ, BRASIL

DIVERSIDAD DE HORMIGAS (HYMENOPTERA: FORMICIDAE) PRESENTES EN AMAPÁ, BRASIL

Monique Telcia dos Santos Damasceno¹, Manoel Daltro Nunes Garcia Junior²

e747692

<https://doi.org/10.47820/recima21.v7i4.7692>

PUBLISHED: 04/2026

ABSTRACT

The order Hymenoptera is one of the megadiverse groups in the class Insecta. Among the members of this order are the ants, all belonging to the family Formicidae. These insects are eusocial and play important roles in ecosystems. The Brazilian myrmecofauna is the largest in the Neotropical region, with approximately 1,700 registered species distributed across 117 genera. However, information on ant diversity is still scarce in different states of the country. Thus, the objective of this study is to present a list of species occurring in the state of Amapá. This list of species from Amapá was compiled based on an analysis of literature contained in databases. The following keywords were used to obtain information: ants, formigas, Neotropical, Brazil, Brasil, Amapá, entomology, entomologia, diversity, diversidade and myrmecofauna. Records contained on specialized websites were also evaluated. This study identified eight subfamilies, 69 genera, and 205 species of ants in the state of Amapá. The most representative subfamilies were Myrmicinae with 27 genera and 88 species, and Ponerinae with 16 genera and 35 species. The genera *Pseudomyrmex* Lund, 1831 (n=17), *Camponotus* Mayr, 1861 (n=16), *Crematogaster* Lund, 1831 (n=13) and *Neoponera* Emery, 1901 (n=13) were those that presented the greatest diversity of species. Although the results of this study show high diversity for the state, further research and expansion of collection areas will certainly increase knowledge of the myrmecofauna in Amapá.

KEYWORDS: Myrmecofauna. Inventory. Distribution. Amazon.

RESUMO

A ordem Hymenoptera é um dos grupos megadiversos da classe Insecta. Entre os integrantes dessa ordem estão as formigas, todas pertencentes à família Formicidae. Esses insetos são eussociais e desempenham funções importantes nos ecossistemas. A mirmecofauna brasileira é a maior da região Neotropical e possui cerca de 1.700 espécies registradas, distribuídas em 117 gêneros. Contudo, em diferentes estados do país, ainda são escassas as informações sobre a diversidade de formigas. Assim, o objetivo do presente estudo é apresentar uma lista das espécies ocorrentes no estado do Amapá. Essa lista de espécies amapaenses foi elaborada a partir da análise de literatura contida em bases de dados. Para a obtenção das informações, foram utilizadas as palavras-chave: ants, formigas, Neotropical, Brazil, Brasil, Amapá, entomology, entomologia, diversity, diversidade e mirmecofauna. Também foram avaliados os registros contidos em sites especializados. O presente estudo apontou a ocorrência de oito subfamílias, 69 gêneros e 205 espécies de formigas no estado do Amapá. As subfamílias mais representativas foram Myrmicinae, com 27 gêneros e 88 espécies, e Ponerinae, com 16 gêneros e 35 espécies. Os gêneros *Pseudomyrmex* Lund, 1831 (n=17), *Camponotus* Mayr, 1861 (n=16), *Crematogaster* Lund, 1831 (n=13) e *Neoponera* Emery, 1901 (n=13) foram os que apresentaram maior diversidade de

¹ Biologist and Master's degree holder in Tropical Biodiversity from the Federal University of Amapá. Conducting entomological and cytogenetic studies with insects of the Formicidae family.

² Biologist, PhD in Tropical Biodiversity. Primarily working on entomological studies in the Amazon region.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

espécies. Embora os resultados deste estudo demonstrem elevada diversidade para o estado, a realização de novas pesquisas e a ampliação das áreas de coleta certamente aumentarão o conhecimento sobre a mirmecofauna do Amapá.

PALAVRAS-CHAVE: Mirmecofauna. Inventário. Distribuição. Amazônia.

RESUMEN

Hymenoptera es uno de los grupos megadiversos de la clase Insecta. Entre los miembros de este orden se encuentran las hormigas, todas pertenecientes a la familia Formicidae. Estos insectos son eusociales y desempeñan funciones importantes en los ecosistemas. La mirmecofauna brasileña es la más grande de la región neotropical, con aproximadamente 1.700 especies registradas pertenecientes a 117 géneros. Sin embargo, en diferentes estados del país aún hay poca información sobre la diversidad de hormigas. Por lo tanto, el objetivo del presente estudio es presentar una lista de especies presentes en el estado de Amapá. Esta lista de especies de Amapá se elaboró a partir del análisis de la bibliografía contenida en bases de datos. Para obtener la información se utilizaron las siguientes palabras clave: ants, formigas, Neotropical, Brazil, Brasil, Amapá, entomology, entomologia, diversity, diversidade y mirmecofauna. También se evaluaron los registros contenidos en sitios web especializados. Este estudio señaló la presencia de ocho subfamilias, 69 géneros y 205 especies de hormigas en el estado de Amapá. Las subfamilias más representativas fueron Myrmicinae, con 27 géneros y 88 especies, y Ponerinae, con 16 géneros y 35 especies. Los géneros *Pseudomyrmex* Lund, 1831 (n=17), *Camponotus* Mayr, 1861 (n=16), *Crematogaster* Lund, 1831 (n=13) y *Neoponera* Emery, 1901 (n=13) fueron los que presentaron mayor diversidad de especies. Aunque los resultados de este estudio muestran una gran diversidad para el estado, la realización de nuevas investigaciones y la ampliación de las áreas de recolección sin duda aumentarán el conocimiento sobre la mirmecofauna en Amapá.

PALABRAS CLAVE: Mirmecofauna. Inventario. Distribución. Amazonia.

INTRODUCTION

Ants belong to the order Hymenoptera and are included in a single family known as Formicidae, within the class Insecta. This is one of the orders with the highest number of species, constituting one of the most megadiverse groups (Huber, 2017; Fernández, 2022a). According to Baccaro *et al.* (2015), approximately 3,000 species of ants occur in the Neotropical Region, included in 13 subfamilies and 142 genera, many of which are endemic. Brazil is home to more than half of the species described for the Neotropical Region: approximately 1,700 species and 117 genera have been recorded in the country (Feitosa; Dias, 2024).

Ants can be found in virtually all terrestrial ecosystems, with the exception of polar regions, some oceanic islands, and high-altitude environments (Ward, 2006). Among terrestrial organisms, this group certainly stands out in terms of diversity, relative abundance, and animal biomass (Kaminski *et al.*, 2009). For example, in the Amazon rainforest, the combined biomass of ants can be up to four times greater than the combined biomass of all vertebrates (Fitkau, 1973). Common characteristics of ants, such as eusocial behavior, overlapping generations, and the enormous variety of intraspecific mechanisms of recognition and chemical communication, are cited as responsible for their impressive numbers in terrestrial environments (Hölldobler; Wilson, 1990).

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

In the Brazilian Amazon, studies conducted on myrmecofauna by (Marques *et al.*, 2002; Peixoto *et al.*, 2010; Vasconcelos *et al.*, 2010; Vieira *et al.*, 2013; Vicente *et al.*, 2016; Schmidt *et al.*, 2020; De Albuquerque *et al.*, 2021; Dias; Feitosa, 2024 Silv;a *et al.*, 2025) have provided relevant information about the group. However, knowledge about ant diversity is still limited for some states in the Amazon.

Amapá is among the states in the Amazon region of Brazil where research on ants is rare. For the state, the studies carried out by Kempf, (1959; 1960a; Oliveira *et al.*, 1995) and Nascimento *et al.*, (2019) certainly deserve mention, due to the number of species recorded. This is because most species records for Amapá are sporadic and scattered across different publications, which makes it difficult to gather information and understand diversity. Thus, the objective of this study is to present a list of ant species found in the state of Amapá.

METHODOLOGY

To compile the list of species occurring in the state, scientific literature containing information on the myrmecofauna of Amapá was investigated. For this purpose, the databases Google Scholar, Science Direct, Scielo, Web of Science, and Scopus were used, as well as specialized ant websites such as AntWeb. The searches were conducted using a series of keywords: ants, formigas, Neotropical, Brazil, Brasil, Amapá, entomology, entomologia, diversity, diversidade and myrmecofauna.

To create the table of species, present in the state, studies found in databases published since the 1950s that mentioned ant species collected in Amapá were evaluated. However, priority was given to studies that presented identification at the species level, the incorporation of organisms at the genus level occurred in cases where there are still no identified species for the state. The validation of the specimens was carried out using data available in recent taxonomic revisions of genera and species of ants present in Brazil.

RESULTS

In total, 205 ant species were identified for Amapá, belonging to 69 genera and eight subfamilies (Table 1). Among the Formicidae subfamilies occurring in the state, Myrmicinae (n=27), Ponerinae (n=16), and Dolichoderinae (n=8) are those with the largest number of genera. The most representative subfamilies in this study were Myrmicinae (n=88) with almost 43% of the identified species, Ponerinae (n=35) with 17%, and Formicinae (n=21) with close to 10% of the species. At the other extreme, Paraponerinae is represented by a single species in the state.

Among the genera identified, *Pseudomyrmex* Lund, 1831 (n=17), *Camponotus* Mayr, 1861 (n=16), *Crematogaster* Lund, 1831 (n=13) and *Neoponera* Emery, 1901 (n=13) were those that presented the greatest number of species. Of the total number of genera occurring in the state, nine



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltro Nunes Garcia Junior

have no recognized species. Ponerinae (n=3) is the subfamily with the largest number of identifications at the generic level only.

DISCUSSION

The state of Amapá is located in the northern region of the country, within the Brazilian Amazon. According to Newbold *et al.*, (2019), this biome comprises about two-thirds of terrestrial biodiversity. This high biodiversity allows for numerous studies, including those on myrmecofauna, to have high numbers of species. For example, for Amazonas the diversity of species/morphotypes reported by Vasconcelos *et al.* (2010) was 166, while Da Silva *et al.*, (2025) identified 205, and in Acre, Dias Feitosa (2024) indicate in their work the existence of 171 species. In Pará, studies by Vicente *et al.*, (2016) and Bochenski *et al.*, (2025) recorded 203 and 237 species, respectively.

Although Amapá is located within the Amazon biome, the ant fauna recognized for the state is still limited; the total of 205 species occurring in Amapá territory represents just over 23% of the Brazilian fauna, which comprises approximately 1,700 species according to Feitosa and Dias (2024). When we compare the number of species recorded for Amapá with that of other states such as Pará (n=753) (De Albuquerque *et al.*, 2021) and Acre (n=394) (Schmidt *et al.*, 2025), for example, it becomes clear that knowledge of the fauna is still very fragile for the state.

In the state, the occurrence of 69 genera was confirmed, a number that is equivalent to approximately 60% of the 117 genera existing in Brazil (Feitosa; Dias, 2024). This number of genera recorded for the state is close to the diversity found in the state of Pará (n=90) (De Albuquerque *et al.*, 2021) and in French Guiana (n=84) (Franco *et al.*, 2019), which are located south and north of the state, respectively.

Among the genera occurring in Amapá, the records of *Iridomyrmex* Mayr, 1862 and *Technomyrmex* Mayr, 1872, both initially made by Oliveira *et al.*, (1995), are considered important for the Brazilian myrmecofauna; these exotic ants have low numbers of records for the country. According to Shattuck (1992), *Iridomyrmex* Mayr, 1862 is a group of ants found in Australia and Asia, while *Technomyrmex* Mayr, 1872 has most of its species recorded in Africa, parts of Asia, and Australia.

The first record of the species *Technomyrmex vitiensis* Mann, 1921 for Brazil was made from specimens collected in the northern region of Amapá, in the municipality of Oiapoque (Barros *et al.*, 2022). *T. vitiensis* Mann, 1921 is a species native to Asia (Väänänen *et al.*, 2018), and its occurrence in Amapá is probably a consequence of the flow of people between French Guiana and the municipality of Oiapoque in Brazil, because during this transit, the transport of various products, including fauna, is common (Almeida; Rauber, 2017; Barros *et al.*, 2022).

Research on ant fauna in Amapá has not been carried out by study groups specializing in Formicidae. This fact can be observed in recent publications for the state, which worked with the



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

cytogenetics of the group (Aguiar *et al.*, 2020; Barros *et al.*, 2021a; 2021b; 2022; Damasceno *et al.*, 2024; Lod *et al.*, 2026) and the transport of bacteria in hospital environments by ants (Nascimento, 2019).

The studies that contributed the most information about the ant fauna of Amapá were carried out by Kempf in the 1960s and 1970s and Oliveira *et al.*, (1995). The last consistent species survey in Amapá was conducted just over 30 years ago. The study by Oliveira *et al.*, (1995) was carried out in the southwestern region of the state and evaluated the ant fauna in eucalyptus plantations. In total, the study presented the occurrence of 121 morphospecies of ants, belonging to six subfamilies and 50 genera.

Although they represent the most significant works for the state, the studies carried out by Kempf in the 1960s and 1970s and Oliveira *et al.*, (1995) present species lists that show a number of inconsistencies with the current taxonomic classification, due to the time lapse since their execution.

CONSIDERATIONS

Although the results of this study demonstrated significant diversity for the state, the geographical characteristics and preserved environmental conditions of Amapá likely guarantee the existence of a much larger number of species. And Amapá will likely solidify its position among the Brazilian states with the greatest ant biodiversity. This confirmation depends on conducting further studies and expanding the collection areas.

Table 1. List of subfamilies, genera and species of ants with recorded occurrences in Amapá

Subfamily / Genus	Species	Reference
Myrmicinae		
<i>Acromyrmex</i> Mayr, 1865		
	<i>A. balzani</i> (Emery, 1890)	Rando and Forti, 2005; Jordão and Silva, 2006; Vinha, 2007
	<i>A. coronatus</i> (Fabricius, 1804)	Kempf, 1972
	<i>A. landolti</i> (Forel, 1885)	Vinha, 2007
	<i>A. laticeps</i> (Emery, 1905)	Delabie <i>et al.</i> , 2011
	<i>A. octospinosus</i> (Reich, 1793)	Céline <i>et al.</i> , 2012
	<i>A. subterraneus subterraneus</i> (Forel, 1893)	Rando and Forti, 2005
<i>Allomerus</i> Mayr, 1878		
	<i>A. decemarticulatus</i> Mayr, 1878	AntWeb
<i>Atta</i> Fabricius, 1804		
	<i>A. cephalotes</i> (Linnaeus, 1758)	Delabie <i>et al.</i> , 2011; Teixeira <i>et al.</i> , 2022

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrio Nunes Garcia Junior

	<i>A. laevigata</i> (Smith 1858)	Vinha, 2007
	<i>A. sexdens</i> Linnaeus, 1758	Jordão and Silva, 2006; Aguiar et al., 2020
	<i>A. sexdens rubropilosa</i> Forel, 1908	Jordão and Silva, 2006
<i>Apterostigma</i> Mayr, 1865		
	<i>A. andense</i> Lattke, 1997	Lod, 2026
	<i>A. jubatum</i> Wheeler, 1925	Lod, 2026
	<i>A. tropicoxa</i> Lattke, 1997	Lod, 2026
<i>Basiceros</i> Schulz, 1906		
	<i>B. tumucumaquensis</i> Probst & Brandão, 2022	AntWeb
	<i>B. scambognathus</i> (Brown, 1949)	AntWeb
<i>Cardiocondyla</i> Emery, 1869		
	<i>C. minutior</i> Forel, 1899	AntWeb
<i>Carebara</i> Westwood, 1840		
	<i>C. urichi</i> (Wheeler, 1922)	Fernández, 2004
<i>Cephalotes</i> Latreille, 1802		
	<i>C. atratus</i> (Linnaeus, 1758)	Oliveira et al., 2021
	<i>C. clypeatus</i> (Fabricius, 1804)	Oliveira et al., 2021
	<i>C. conspersus</i> (Smith, 1867)	Oliveira et al., 2021
	<i>C. cordatus</i> (Smith, 1853)	Oliveira et al., 2021; Damasceno et al., 2024
	<i>C. depressus</i> (Klug, 1824)	Oliveira et al., 2021
	<i>C. maculatus</i> (Smith, 1876)	Oliveira et al., 2021
	<i>C. minutus</i> (Fabricius, 1804)	Oliveira et al., 2021
	<i>C. oculatus</i> (Spinola, 1851)	Oliveira et al., 2021
	<i>C. opacus</i> Santschi, 1920	Oliveira et al., 2021
	<i>C. pavonii</i> (Latreille, 1809)	Oliveira et al., 2021
	<i>C. placidus</i> (Smith, 1860)	Oliveira et al., 2021
	<i>C. pusillus</i> (Klug, 1824)	Oliveira et al., 2021
	<i>C. serratriceps</i> (Smith, 1858)	Oliveira et al., 2021
<i>Crematogaster</i> Lund, 1831		
	<i>C. abstinens</i> Forel, 1899	AntWeb
	<i>C. amapaensis</i> Kempf, 1960	Kempf, 1960a
	<i>C. brasiliensis</i> Mayr, 1878	Felizardo, 2010
	<i>C. boliviana</i> (Wheeler, 1922)	Kempf, 1960b
	<i>C. crinosa</i> Mayr, 1862	Kempf, 1972
	<i>C. erecta</i> (Mayr, 1866)	Silveira, 2022
	<i>C. levior</i> Longino, 2003	Felizardo, 2010
	<i>C. limata</i> (Smith, 1858)	Silveira, 2022
	<i>C. longispina</i> Emery, 1890	Felizardo, 2010
	<i>C. obscurata</i> Emery, 1895	Kempf, 1972
	<i>C. stollii</i> Forel, 1885	AntWeb
	<i>C. sotobosque</i> Longino, 2003	AntWeb
	<i>C. tenuicula</i> (Forel, 1904)	Silveira, 2022
<i>Cyphomyrmex</i> Mayr, 1862		
	<i>C. laevigatus</i> Weber, 1938	Damasceno et al., 2024
	<i>C. rimosus</i> (Spinola, 1851)	Kempf, 1972
	<i>C. transversus</i> Emery, 1894	Lod, 2026
<i>Daceton</i> Perty, 1833		

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

	<i>D. armigerum</i> (Latreille, 1802)	Kempf, 1972
<i>Lachnomyrmex</i> Wheeler, 1910		
	<i>L. pilosus</i> Weber, 1950	Feitosa and Brandao, 2008
<i>Megalomyrmex</i> Forel, 1885		
	<i>M. bituberculatus</i> (Fabricius, 1798)	Kempf, 1972
	<i>M. incisus</i> Smith, 1947	Ferreira et al., 2023; Damasceno et al., 2024
	<i>M. leoninus</i> Forel, 1885	AntWeb
<i>Monomorium</i> Mayr, 1855		
	<i>M. floricola</i> (Forel, 1893)	Sá, 2013; Nascimento, 2019
	<i>M. pharaonis</i> (Mayr, 1855)	Sá, 2013; Nascimento, 2019
<i>Mycetophylax</i> Emery, 1913		
	<i>M. bigibbosus</i> (Emery, 1894)	Kempf, 1972
<i>Myrmicocrypta</i> Smith, 1860		
	<i>M. bucki</i> Sosa-Calvo & Schultz, 2010	Sosa-Calvo and Schultz, 2010
<i>Nesomyrmex</i> Wheeler, 1910		
	<i>N. pleuriticus</i> (Kempf, 1959)	Kempf, 1960a; 1972
	<i>N. spininodis</i> (Mayr, 1887)	AntWeb
<i>Ochetomyrmex</i> Mayr, 1878		
	<i>O. semipolitus</i> Mayr, 1878	Kempf, 1972
<i>Octostruma</i> Forel, 1912		
	<i>Octostruma</i> sp.	Oliveira et al., 1995
<i>Pheidole</i> Westwood, 1839		
	<i>P. biconstricta</i> Mayr, 1870	Kempf, 1972
	<i>P. fallax</i> Maio, 1870	Kempf, 1972
	<i>P. jelskii</i> Mayr, 1884	Damasceno et al., 2024
	<i>P. mamore</i> Mann, 1916	AntWeb
	<i>P. radoszkowskii</i> Mayr, 1884	Kempf, 1972
	<i>P. vorax</i> (Fabricius, 1804)	Damasceno et al., 2024
<i>Pogonomyrmex</i> Mayr, 1868		
	<i>P. naegelii</i> Emery, 1878	AntWeb
<i>Procryptocerus</i> Emery, 1887		
	<i>P. attenuatus</i> (Smith, 1876)	AntWeb
	<i>P. hylaeus</i> Kempf, 1951	Leitão et al., 2025
	<i>P. paleatus</i> Emery, 1896	Kempf, 1960a
	<i>P. subpilosus</i> (Smith, 1860)	AntWeb
<i>Sericomyrmex</i> Mayr, 1865		
	<i>S. mayri</i> Forel, 1912	AntWeb
<i>Solenopsis</i> Westwood, 1840		
	<i>S. altinodis</i> Forel, 1912	AntWeb
	<i>S. saevissima</i> (Smith, 1855)	Reis and Souto, 2011; Sá, 2013; Nascimento, 2019
	<i>S. geminata</i> (Fabricius, 1804)	Martins and Souto, 2021
	<i>S. globularia</i> (Smith, 1858)	Sá, 2013
<i>Strumigenys</i> Smith, 1860		
	<i>S. ascita</i> Bolton, 2000	Bolton, 2000
	<i>S. cordovensis</i> Mayr, 1887	Kempf, 1972

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

	<i>S. denticulata</i> Mayr, 1887	INPA
	<i>S. elongata</i> Roger, 1863	INPA
	<i>S. louisianae</i> Roger, 1863	Barros et al., 2021a
	<i>S. perparva</i> Brown, 1958	Bolton, 2000
	<i>S. schulzi</i> Emery, 1894	Damasceno et al., 2024
	<i>S. subdentata</i> Mayr, 1887	De Albuquerque et al., 2021
	<i>S. stenotes</i> (Bolton, 2000)	Bolton, 2000
	<i>S. villiersi</i> (Perrault, 1986)	Antweb
	<i>S. vilhenai</i> Bolton, 2000	Antweb
<i>Trachymyrmex</i> Forel, 1893		
	<i>Trachymyrmex</i> sp.	Oliveira et al., 1995
<i>Wasmannia</i> Forel, 1893		
	<i>W. auropuctata</i> (Roger, 1863)	Nascimento, 2019
	<i>W. scrobifera</i> Kempf, 1961	INPA
Paraponerinae		
<i>Paraponera</i> Smith, 1858		
	<i>P. clavata</i> (Fabricius, 1775)	Kempf, 1972
Ponerinae		
<i>Anochetus</i> Mayr, 1861		
	<i>A. bispinosus</i> (Smith, 1858)	Kempf, 1972
	<i>A. neglectus</i> Emery, 1894	INPA
<i>Centromyrmex</i> Mayr, 1866		
	<i>C. brachycola</i> (Roger, 1861)	AntWeb
	<i>C. gigas</i> Forel, 1911	AntWeb
<i>Cryptopone</i> Emery, 1893		
	<i>C. guianensis</i> (Weber, 1939)	AntWeb
<i>Discothyrea</i> Roger, 1863		
	<i>Discothyrea</i> sp	Oliveira et al., 1995
<i>Ectatomma</i> Smith, 1858		
	<i>Ectatomma</i> sp.	Oliveira et al., 1995
<i>Gnamptogenys</i> Roger, 1863		
	<i>G. lanei</i> Kempf, 1960	Kempf, 1960a
<i>Heteroponera</i> Mayr, 1887		
	<i>Heteroponera</i> sp.	Oliveira et al., 1995; Reis and Souto, 2011
<i>Hypoponera</i> Santschi, 1938		
	<i>H. opaciceps</i> (Mayr, 1887)	Kempf, 1972
<i>Leptogenys</i> Roger, 1861		
	<i>L. unistimulosa</i> Roger, 1863	Kempf, 1972
<i>Mayaponera</i> Schmidt & Shattuck, 2014		
	<i>M. constricta</i> (Mayr, 1884)	AntWeb
<i>Neoponera</i> Emery, 1901		
	<i>N. apicalis</i> (Latreille, 1802)	Kempf, 1972; MacKay and MacKay, 2010
	<i>N. bactronica</i> (Fernandes, De Oliveira & Delabie, 2014)	Fernandes et al., 2014
	<i>N. commutata</i> (Roger, 1860)	Kempf, 1972; Troya and Lattke, 2022
	<i>N. crenata</i> (Roger, 1861)	Kempf, 1972; MacKay and MacKay, 2010
	<i>N. curvinodis</i> (Forel, 1899)	Fernandes et al., 2014

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

	<i>N. foetida</i> (Linnaeus, 1758)	Kempf, 1972
	<i>N. inversa</i> (Smith, 1858)	AntWeb
	<i>N. mashpi</i> Troya & Lattke, 2022	Troya and Lattke, 2022
	<i>N. oberthueri</i> (Emery, 1890)	Kempf, 1972; MacKay and MacKay, 2010
	<i>N. obscuricornis</i> (Emery, 1890)	Kempf, 1972
	<i>N. unidentata</i> Mayr, 1862	Damasceno et al., 2024
	<i>N. striatinodis</i> (Emery, 1890)	AntWeb
	<i>N. villosa</i> (Fabricius, 1804)	AntWeb
<i>Odontomachus</i> Latreille, 1804		
	<i>O. bauri</i> Emery, 1892	AntWeb
	<i>O. affinis</i> Guérin-Méneville, 1844	Kempf, 1959
	<i>O. caelatus</i> Brown, 1976	AntWeb
	<i>O. haematodus</i> (Linnaeus, 1758)	Kempf, 1959; 1972
	<i>O. hastatus</i> (Fabricius, 1804)	Kempf, 1960a; 1972
	<i>O. laticeps</i> Roger, 1861	Kempf, 1972
	<i>O. panamensis</i> Forel, 1899	Brown, 1976
	<i>O. opaciventris</i> Forel, 1899	Kempf, 1959
<i>Pachycondyla</i> Smith, 1858		
	<i>P. crassinoda</i> (Latreille, 1802)	AntWeb
	<i>P. harpax</i> (Fabricius, 1804)	AntWeb
<i>Platythyrea</i> Roger, 1863		
	<i>P. sinuata</i> (Roger, 1860)	Kempf, 1972; Brandao, 1991
<i>Pseudoponera</i> Emery, 1900		
	<i>P. stigma</i> (Fabricius, 1804)	Damasceno et al., 2024
<i>Wadeura</i> Weber, 1939		
	<i>W. guianensis</i> Weber, 1939	MacKay and MacKay, 2010
Formicinae		
<i>Brachymyrmex</i> Mayr, 1868		
	<i>B. coactus</i> Mayr, 1887	AntWeb
	<i>B. heeri</i> Forel, 1874	Damasceno et al., 2024
<i>Camponotus</i> Mayr, 1861		
	<i>C. atriceps</i> (Smith, 1858)	Kempf, 1959; 1972; Hashmi, 1973
	<i>C. blandus</i> (Smith, 1858)	AntWeb
	<i>C. coloratus</i> Forel, 1904	Santos and Sobrinho, 2025
	<i>C. chartifex</i> (Smith, 1860)	Fernández, 2002
	<i>C. coruscus</i> (Smith, 1862)	AntWeb
	<i>C. eurynotus</i> Forel, 1907	Kempf, 1960a; 1972
	<i>C. femoratus</i> (Fabricius, 1804)	Kempf, 1959; 1972
	<i>C. latangulus</i> Roger, 1863	Kempf, 1959; 1972
	<i>C. lenkoi</i> Kempf, 1960	Kempf, 1960a
	<i>C. leydigi</i> Forel, 1886	Kempf, 1960a; 1972
	<i>C. nidulans</i> (Smith, 1860)	Fernández, 2002
	<i>C. rapax</i> (Fabricius, 1804)	AntWeb
	<i>C. senex</i> (Smith, 1858)	AntWeb

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltro Nunes Garcia Junior

	<i>C. sexguttatus</i> (Fabricius, 1793)	Nascimento, 2019
	<i>C. urichi</i> Forel, 1899	Kempf, 1960a; 1972
	<i>C. vittatus</i> (Forel, 1904)	Sá, 2013
<i>Gigantiops</i> Roger, 1863		
	<i>G. destructor</i> (Fabricius, 1804)	Kempf and Lenko, 1968; Kempf, 1972
<i>Nylanderia</i> Emery, 1906		
	<i>Nylanderia</i> sp	Sá, 2013
<i>Paratrechina</i> Motschoulsky, 1863		
	<i>P. longicornis</i> (Latreille, 1802)	Sá, 2013; Nascimento, 2019
Pseudomyrmecinae		
<i>Pseudomyrmex</i> Lund, 1831		
	<i>P. browni</i> Kempf, 1967	Kempf, 1967; Brandão et al., 2010
	<i>P. concolor</i> (Smith, 1860)	Pacheco Junior, 2016; Pacheco Junior and Del-Claro, 2021
	<i>P. filiformis</i> (Fabricius, 1804)	Kempf, 1960a
	<i>P. godmani</i> (Forel, 1899)	Kempf, 1967; Brandão et al., 2010
	<i>P. gracilis</i> (Fabricius, 1804)	Kempf, 1959; 1972
	<i>P. laevifrons</i> Ward, 1989	Ward, 2017
	<i>P. malignus</i> (Wheeler, 1921)	Ward, 1999
	<i>P. penetrator</i> (Smith, 1877)	Ward, 1999
	<i>P. pupa</i> (Forel, 1911)	Kempf, 1960a; 1972
	<i>P. subater</i> (Wheeler & Mann, 1914)	AntWeb
	<i>P. simplex</i> (Smith, 1877)	AntWeb
	<i>P. tenuis</i> (Fabricius, 1804)	Kempf, 1972
	<i>P. termitarius</i> (Smith, 1855)	Kempf, 1960a
	<i>P. triplaridis</i> (Forel, 1904)	Ward, 1999
	<i>P. unicolor</i> (Smith, 1855)	Kempf, 1960a; 1972
	<i>P. viduus</i> (Smith, 1858)	Ward, 1999
	<i>P. vinneri</i> (Forel, 1906)	AntWeb
Dolichoderinae		
<i>Azteca</i> Forel, 1878		
	<i>Azteca</i> sp.	Oliveira et al., 1995
<i>Dorymyrmex</i> Mayr, 1866		
	<i>D. goeldii</i> Forel, 1904	AntWeb
	<i>D. brunneus</i> Forel, 1908	AntWeb
<i>Dolichoderus</i> Lund, 1831		
	<i>D. attelaboides</i> (Fabricius, 1775)	Kempf, 1959; 1972
	<i>D. bidens</i> (Linnaeus, 1758)	Barros et al., 2021b
	<i>D. bispinosus</i> (Olivier, 1792)	Kempf, 1959; 1972
	<i>D. decollatus</i> Smith, 1858	Kempf, 1959; 1972
	<i>D. fernandesi</i> Mackay, 1993	Mackay, 1993
	<i>D. ferrugineus</i> Forel, 1903	Kempf, 1960a; 1972
	<i>D. gagates</i> Emery, 1890	Kempf, 1960a; 1972; Mackay, 1993
	<i>D. imitator</i> Emery, 1894	AntWeb
	<i>D. lutosus</i> (Smith, 1858)	Kempf, 1972

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

	<i>D. quadridentatus</i> (Roger, 1862)	Kempf, 1960a; 1972
	<i>D. septemspinus</i> Emery, 1894	Kempf, 1960a; 1972
	<i>D. tristis</i> Mann, 1916	Mackay, 1993
<i>Gracilidris</i> Wild & Cuzzo, 2006		
	<i>G. pombero</i> Wild & Cuzzo, 2006	AntWeb
<i>Iridomyrmex</i> Mayr, 1862		
	<i>Iridomyrmex</i> sp.	Oliveira et al., 1995
<i>Linepithema</i> Mayr, 1866		
	<i>L. humile</i> (Mayr, 1868)	Nascimento, 2019
<i>Tapinoma</i> Foerster, 1850		
	<i>T. melanocephalum</i> Fabricius, 1793	Nascimento, 2019; Teixeira et al., 2025
<i>Technomyrmex</i> Mayr, 1872		
	<i>T. vitiensis</i> Mann, 1921	Barros et al., 2022
Dorylinae		
<i>Acanthostichus</i> Mayr, 1887		
	<i>A. femoralis</i> Kusnezov, 1962	AntWeb
	<i>A. fuscipennis</i> Emery, 1895	MacKay, 1996
	<i>A. quadratus</i> Emery, 1895	AntWeb
<i>Cheliomyrmex</i> Mayr, 1870		
	<i>C. morosus</i> (Smith, 1859)	Kempf, 1960a
<i>Eciton</i> Latreille, 1804		
	<i>E. burchellii</i> (Westwood, 1842)	Kempf, 1959
	<i>E. drepanophorum</i> Smith, 1858	Kempf, 1959; 1972
	<i>E. hamatum</i> (Fabricius, 1782)	Kempf, 1959; 1972
	<i>E. setigaster</i> Borgmeier, 1953	Kempf, 1959; 1972
	<i>E. vagans</i> (Olivier, 1792)	Kempf, 1972; Watkins, 1976
<i>Labidus</i> Jurine, 1807		
	<i>L. coecus</i> (Latreille, 1802)	Kempf, 1960a; 1972; Watkins, 1976
	<i>L. praedator</i> (Smith, 1858)	Kempf, 1959; 1972; Watkins, 1976
	<i>L. truncatidens</i> (Santschi, 1920)	Kempf, 1960a; 1972; Watkins, 1976
<i>Neivamyrmex</i> Borgmeier, 1940		
	<i>N. emersoni</i> (Wheeler, 1921)	AntWeb
	<i>N. falcifer</i> (Emery, 1900)	Kempf, 1959; 1972; Watkins, 1976
	<i>N. halidaii</i> (Shuckard, 1840)	Kempf, 1960a; 1972; Watkins, 1976
	<i>N. walkerii</i> (Westwood, 1842)	Kempf, 1960a; 1972; Watkins, 1976
<i>Nomamyrmex</i> Borgmeier, 1936		
	<i>N. hartigii</i> (Westwood, 1842)	Kempf, 1960a; 1972; Watkins, 1976

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

Ectatomminae		
<i>Ectatomma</i> Smith, 1858		
	<i>E. brunneum</i> Smith, 1858	Kempf, 1959; 1972
	<i>E. lugens</i> Emery, 1894	Kempf, 1959; 1972
	<i>E. tuberculatum</i> (Olivier, 1792)	Kempf, 1959; 1972
<i>Gnamptogenys</i> Roger, 1863		
	<i>G. annulata</i> (Mayr, 1887)	Kempf, 1960a; 1972; Camacho et al., 2020
	<i>G. concinna</i> (Smith, 1858)	Delabie et al., 2010
<i>Heteroponera</i> Mayr, 1887		
	<i>H. dentinodis</i> (Mayr, 1887)	Kempf, 1972; Feitosa, 2011
<i>Holcoponera</i> Mayr, 1887		
	<i>H. pleurodon</i> (Emery, 1896)	Kempf, 1960a; 1972
<i>Poneracantha</i> Emery, 1897		
	<i>P. lanei</i> (Kempf, 1960)	Kempf, 1972; Camacho et al., 2020
	<i>P. triangularis</i> (Mayr, 1887)	AntWeb

REFERENCES

ALMEIDA, C. S.; RAUBER, A. L. Oiapoque, aqui começa o Brasil: a fronteira em construção e os desafios do desenvolvimento regional. **Redes**, v. 22, n. 1, p. 474–493, 2017. DOI: <https://doi.org/10.17058/redes.v22i1.8532>.

ANTWEB. **California Academy of Science**. Disponível em: <https://www.qbif.org/dataset/13b70480-bd69-11dd-b15f-b8a03c50a862>. Acesso em: jan. 2026.

BACCARO, F. B. *et al.* **Guia para os gêneros de formigas do Brasil**. Manaus: Editora INPA, 2015. 388 p.

BARROS, L. A. C. *et al.* Cytogenetic survey of eight ant species from the Amazon rainforest. **Caryologia**, v. 74, n. 4, p. 29–38, 2021b. DOI: <https://doi.org/10.36253/caryologia-11061>.

BARROS, L. A. C. *et al.* Cytogenetics of *Strumigenys louisianae* Roger, 1863 (Formicidae: Myrmicinae) from North eastern Amazonia shed light on a difficult species complex. **Zoologischer Anzeiger**, v. 294, p. 100–105, 2021a.

BARROS, L. A. C. *et al.* First report of the tramp ant *Technomyrmex vitiensis* Mann, 1921 (Formicidae: Dolichoderinae) in Brazil with cytogenetic and sperm structure data and an updated key to Brazilian Dolichoderinae genera. **Zoological Studies**, v. 61, e29, 2022. DOI: <https://doi.org/10.6620/zs.2022.61-29>.

BOCHENSKI, W. S. *et al.* Ant brightness and size influence the seasonal vegetation use by ground-dwelling ants (Hymenoptera: Formicidae) in an Amazonia-Cerrado transition zone. **Ecological Entomology**, v. 50, n. 6, p. 1082–1095, 2025. DOI: <https://doi.org/10.1111/een.70000>.

BOLTON, B. The ant tribe Dacetini. **Memoirs of the American Entomological Institute**, v. 65, p. 1–1028, 2000.

ISSN: 2675-6218 - RECIMA21

This article is published in Open Access under the Creative Commons Attribution 4.0 International (CC-BY) license, which allows unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



BRANDÃO, C. R. F.; ESTEVES, F. A.; PRADO, L. A. A catalogue of the Pseudomyrmecinae ants type specimens (Hymenoptera, Formicidae) deposited in the Museu de Zoologia da Universidade de São Paulo, Brazil. **Papéis Avulsos de Zoologia**, v. 50, n. 45, p. 693–699, 2010.

BROWN, W. L. Jr. Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section A. Introduction, subtribal characters. Genus **Odontomachus**. **Studia Entomologica**, v. 19, p. 67–171, 1976.

CAMACHO, G. P.; FRANCO, W.; FEITOSA, R. M. Additions to the taxonomy of *Gnamptogenys* Roger (Hymenoptera: Formicidae: Ectatomminae) with an updated key to the New World species. **Zootaxa**, v. 4747, p. 450–476, 2020.

CÉLINI, L. L. *et al.* Présence et origine d'*Acromyrmex octospinosus* (Reich, 1793) à Saint-Barthélemy, Petites Antilles (Hymenoptera, Formicidae, Attini). **Bulletin de la Société entomologique de France**, v. 117, n. 2, p. 167–172, 2012.

DA SILVA, C. V. C. *et al.* The structure of ant assemblages on Amazonian fluvial islands: the roles of recurrent disturbance and isolation. **Oecologia**, v. 207, n. 8, p. 140, 2025.

DAMASCENO, M. T. S. *et al.* Physical chromosomal mapping of major ribosomal genes in 15 ant species with a review of hypotheses regarding evolution of the number and position of NORs in ants. **Comparative Cytogenetics**, v. 18, p. 105–122, 2024. DOI: <https://doi.org/10.3897/compcytogen.18.125235>.

DE AGUIAR, H. J. A. C. *et al.* Cytogenetic data for sixteen ant species from North-eastern Amazonia with phylogenetic insights into three subfamilies. **Comparative Cytogenetics**, v. 14, n. 1, p. 43–60, 2020. DOI: <https://doi.org/10.3897/CompCytogen.v14i1.46692>.

DE ALBUQUERQUE, E. Z. *et al.* Ants of the State of Pará, Brazil: a historical and comprehensive dataset of a key biodiversity hotspot in the Amazon Basin. **Zootaxa**, v. 5001, n. 30, p. 83, 2021.

DELABIE, J. H. C. *et al.* Distribuição das formigas cortadeiras dos gêneros *Acromyrmex* e *Atta* no Novo Mundo. In: DELLA LUCIA, T. M. C. (org.). **As formigas cortadeiras**. Viçosa: Editora da Universidade Federal de Viçosa, 2011. p. 80–101.

DELÁBIE, J. H. C. *et al.* *Gnamptogenys concinna* (F. Smith, 1858): nouvelles données sur sa distribution et commentaires sur ce cas de gigantisme. **Bulletin de la Société entomologique de France**, v. 115, n. 3, p. 269–277, 2010.

DIAS, A. M.; FEITOSA, R. M. Frontier ants (Hymenoptera: Formicidae) unveiled: exploring the remarkable diversity of the Brazilian Far West. **Biota Neotropica**, v. 24, n. 4, e20241686, 2024. DOI: <https://doi.org/10.1590/1676-0611BN-2024-1686>.

FEITOSA, R. M.; BRANDÃO, C. R. F. A taxonomic revision of the Neotropical myrmicine ant genus *Lachnomyrmex* Wheeler (Hymenoptera: Formicidae). **Zootaxa**, v. 1890, p. 1–49, 2008.

FEITOSA, R. M.; DIAS, A. M. An illustrated guide for the identification of ant subfamilies and genera in Brazil. **Insect Systematics & Evolution**, v. 55, n. 5, p. 451–571, 2024. DOI: <https://doi.org/10.1163/1876312X-bja10062>.

FEITOSA, S. M. R. **Revisão taxonômica e análise filogenética de Heteroponerinae (Hymenoptera, Formicidae)**. 2011. 311 f. Tese (Doutorado) – Universidade de São Paulo, São Paulo, 2011.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

- FELIZARDO, S. P. de S. **Revisão taxonômica do grupo *limata* de *Crematogaster* Lund, 1831 (Formicidae: Myrmicinae: Crematogastrini)**. 2010. 74 f. Dissertação (Mestrado) – Museu Paraense Emílio Goeldi; Universidade Federal do Pará, Programa de Pós-Graduação em Zoologia, Belém, 2010.
- FERNANDES, I. O.; OLIVEIRA, M. L.; DELABIE, J. H. C. Description of two new species in the Neotropical *Pachycondyla foetida* complex (Hymenoptera: Formicidae: Ponerinae) and taxonomic notes on the genus. **Myrmecological News**, v. 19, p. 133–163, 2014.
- FERNÁNDEZ, F. On the diversity of Neotropical Hymenoptera – Sobre la diversidad de Hymenoptera neotropicales. **Caldasia**, v. 43, n. 3, p. 502–513, 2022.
- FERNÁNDEZ, F. Revisión de las hormigas *Camponotus* subgénero *Dendromyrmex* (Hymenoptera: Formicidae). **Papéis Avulsos de Zoologia**, v. 42, p. 47–101, 2002.
- FERNÁNDEZ, F. The American species of the myrmicine ant genus *Carebara* Westwood (Hymenoptera: Formicidae). **Caldasia**, v. 26, p. 191–238, 2004.
- FERREIRA, P. C. *et al.* Citogenética clássica da formiga *Megalomyrmex aff. incisus* coletada em Oiapoque, Amapá, Amazônia. In: CONGRESSO BRASILEIRO INTERDISCIPLINAR EM CIÊNCIA E TECNOLOGIA, 2023, Diamantina. **Anais [...]** Diamantina: Even3, 2023. Disponível em: <https://www.even3.com.br/anais/cobicet2023/651625>. Acesso em: jan. 2026.
- FITKAU, E. J. Role of caimans in the nutrient regime of mouth-lakes of Amazon affluents. **Biotropica**, v. 2, n. 2, p. 138–142, 1973.
- FRANCO, W. *et al.* First checklist of the ants (Hymenoptera: Formicidae) of French Guiana. **Zootaxa**, v. 4674, n. 5, p. 509–543, 2019. DOI: <http://doi.org/10.11646/zootaxa.4674.5.2>.
- HASHMI, A. A. A revision of the Neotropical ant subgenus *Myrmothrix* of genus *Camponotus* (Hymenoptera: Formicidae). **Studia Entomologica**, v. 16, n. 1-4, p. 1–140, 1973.
- HÖLDOBLER, B.; WILSON, E. O. **The ants**. Cambridge: Belknap Press/Harvard University Press, 1990. 732 p.
- HUBER, J. T. Biodiversity of Hymenoptera. In: FOOTIT, R. G.; ADLER, P. H. (ed.). **Insect biodiversity: science and society**. 2. ed. Hoboken: John Wiley & Sons, 2017. p. 419–461.
- INSTITUTO NACIONAL DE PESQUISAS DA AMAZÔNIA (INPA). Coleção do museu (GBIF). Disponível em: <https://www.gbif.org/>. Acesso em: dez. 2025.
- JORDÃO, A. L.; SILVA, R. A. **Guia de pragas agrícolas para o manejo integrado no estado do Amapá**. Ribeirão Preto: Holos, 2006. 182 p.
- KAMINSKI, L. A. *et al.* Ecologia comportamental na interface formiga-planta-herbívoros: interações entre formigas e lepidópteros. **Oecologia Brasiliensis**, v. 13, n. 1, p. 27–44, 2009.
- KEMPF, W. W. Catálogo abreviado das formigas da região Neotropical. **Studia Entomologica**, v. 15, p. 3–344, 1972.
- KEMPF, W. W. Estudos sobre *Pseudomyrmex*. IV (Hymenoptera: Formicidae). **Revista Brasileira de Entomologia**, v. 12, p. 1–12, 1967.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Daltrô Nunes Garcia Junior

KEMPF, W. W. Insecta Amapaensia – Hymenoptera: Formicidae (segunda contribuição). **Studia Entomologica**, v. 3, p. 385–400, 1960a.

KEMPF, W. W. Insecta Amapaensia – Hymenoptera: Formicidae. **Studia Entomologica**, v. 2, p. 209–218, 1959.

KEMPF, W. W. *Tranopeltoides* Wheeler, a synonym of *Creमतogaster* Lund (Hymenoptera, Formicidae). **Entomological News**, v. 71, p. 173–175, 1960b.

KEMPF, W. W.; LENKO, K. Novas observações e estudos sobre *Gigantiops destructor* (Fabricius) (Hymenoptera: Formicidae). **Papéis Avulsos de Zoologia**, v. 21, p. 209–230, 1968.

LEITÃO, E. de A.; BARROS, L. A. C.; DE AGUIAR, H. J. C. Citogenética clássica da formiga *Procryptocerus hylaeus* Kempf, 1951. In: CONGRESSO BRASILEIRO INTERDISCIPLINAR EM CIÊNCIA E TECNOLOGIA, 2025, Diamantina. **Anais [...]** Diamantina: Even3, 2025. Disponível em: <https://www.even3.com.br/anais/congresso-brasileiro-interdisciplinar-ciencia-tecnologia/1197336>. Acesso em: jan. 2026.

LOD, R. B. Citogenética molecular em formigas dos gêneros *Cyphomyrmex* Mayr, 1862 e *Apterostigma* Mayr, 1865 (Formicidae: Myrmicinae) do Escudo das Guianas. 2026. 76 f. Dissertação (Mestrado) – Universidade Federal do Amapá, Programa de Pós-Graduação em Biodiversidade Tropical, Macapá, 2026.

MACKAY, W. P. A review of the New World ants of the genus *Dolichoderus* (Hymenoptera: Formicidae). **Sociobiology**, v. 22, n. 1, p. 1–148, 1993.

MACKAY, W. P. A revision of the ant genus *Acanthostichus* (Hymenoptera: Formicidae). **Sociobiology**, v. 27, p. 129–179, 1996.

MACKAY, W. P.; MACKAY, E. E. **The systematics and biology of the New World ants of the genus *Pachycondyla* (Hymenoptera: Formicidae)**. Lewiston: Edwin Mellen Press, 2010. 642 p.

MARQUES, A. P. C.; ALE-ROCHA, R.; RAFAEL, J. S. Levantamento de formigas (Hymenoptera: Formicidae) em residências de Manaus, estado do Amazonas, Brasil. **Acta Amazonica**, v. 32, n. 1, p. 133–140, 2002.

MARTINS, M. J. L.; SOUTO, R. N. P. Revisão de gêneros de formigas (Hymenoptera: Formicidae) coletadas em floresta de várzea na Amazônia oriental. In: **CONGRESSO AMAPAENSE DE INICIAÇÃO CIENTÍFICA**, 10., 2021. Livro de resumos [...] 2021.

NASCIMENTO, L. E. **Formigas (Hymenoptera: Formicidae) como potenciais vetores mecânicos de micro-organismos patogênicos em um hospital**. 2019. 85 f. Dissertação (Mestrado) – Universidade Federal do Amapá, Programa de Pós-Graduação em Ciências da Saúde, 2019.

NEWBOLD, T. *et al.* Climate and land-use change homogenise terrestrial biodiversity, with consequences for ecosystem functioning and human well-being. **Emerging Topics in Life Sciences**, v. 3, p. 207–219, 2019.

OLIVEIRA, A. M.; POWELL, S.; FEITOSA, R. M. A taxonomic study of the Brazilian turtle ants (Formicidae: Myrmicinae: *Cephalotes*). **Revista Brasileira de Entomologia**, v. 65, n. 3, e20210028, 2021.



REVISTA CIENTÍFICA - RECIMA21 ISSN 2675-6218

DIVERSITY OF ANTS (HYMENOPTERA: FORMICIDAE) OCCURRING IN AMAPÁ, BRAZIL
Monique Telcia dos Santos Damasceno, Manoel Dalro Nunes Garcia Junior

OLIVEIRA, M. A. *et al.* A fauna de formigas em povoamentos de eucalipto e mata nativa no estado do Amapá. **Acta Amazonica**, v. 25, n. 1-2, p. 117–126, 1995.

PACHECO JUNIOR, P. S. M. **Ecologia comportamental e de interações entre *Pseudomyrmex concolor* Smith (Hymenoptera, Formicidae, Pseudomyrmecinae) e sua planta hospedeira *Tachigali myrmecophila* Ducke (Fabaceae: Caesalpinioideae) na Amazônia amapaense.** 2016. 156 f. Tese (Doutorado) – Universidade de São Paulo, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, 2016.

PACHECO JUNIOR, P. S. M.; DEL-CLARO, K. Spatio-temporal variation influences the division of labour in *Pseudomyrmex concolor* Smith (Formicidae: Pseudomyrmecinae). **Journal of Ethology**, v. 39, p. 205–215, 2021. DOI: <https://doi.org/10.1007/s10164-021-00695-1>.

PEIXOTO, T. S. *et al.* Composição e riqueza de formigas (Hymenoptera: Formicidae) em savana e ambientes associados de Roraima. **Revista Agro@ambiente On-line**, v. 4, n. 1, p. 1–10, 2010.

RANDO, J. S. S.; FORTI, L. C. Ocorrência de formigas *Acromyrmex* Mayr em alguns municípios do Brasil. **Acta Scientiarum Biological Sciences**, v. 27, n. 2, p. 129–133, 2005.

REIS, N. P. de J.; SOUTO, R. N. P. Ocorrência de *Solenopsis saevissima* Fr. Smith associada à carcaça de suíno em dois ambientes de cerrado no Campus Marco Zero da Universidade Federal do Amapá. In: CONGRESSO DE ECOLOGIA DO BRASIL, 10., 2011, São Lourenço. **Anais [...]** 2011

SÁ, D. M. da C. **Diversidade de Formicidae em ambiente hospitalar público do município de Macapá-Amapá.** 2013. 80 f. Dissertação (Mestrado) – Universidade Federal do Amapá, Programa de Pós-Graduação em Ciências da Saúde, Macapá, 2013.

SANTOS, C. D. A.; SOBRINHO, T. G. Surveying ant fauna in northern Espírito Santo, Brazil: new distribution records from the Coleção Zoológica Norte Capixaba. **Sociobiology**, v. 72, n. 4, e11539, 2025. DOI: <https://doi.org/10.13102/sociobiology.v72i4.11539>.

SCHMIDT, F. A. *et al.* Ant diversity studies in Acre: what we know and what we could do to know more? **Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais**, v. 15, n. 1, p. 113–134, 2020. DOI: <http://doi.org/10.46357/bcnaturais.v15i1.235>.

SCHMIDT, F. A. *et al.* Western frontier in Brazilian ant diversity: contribution of Rede BIA to ant diversity known in southwestern Amazon. In: **Brazilian myrmecology: exploring the world's richest ant fauna.** São Paulo: Editora Científica, 2025. p. 465–478. DOI: <https://dx.doi.org/10.37885/250920263>.

SHATTUCK, S. O. Generic revision of the ant subfamily Dolichoderinae (Hymenoptera: Formicidae). **Sociobiology**, v. 21, p. 1–181, 1992.

SILVA, E. *et al.* What can ground-dwelling ants tell us about different land-use systems in the Brazilian Amazon? **Forests**, v. 16, n. 7, p. 1190, 2025. DOI: <https://doi.org/10.3390/f16071190>.

SILVEIRA, L. I. **Citogenética clássica e molecular em formigas do gênero *Crematogaster* Lund, 1831 (Formicidae: Hymenoptera) com sondas específicas e de DNA repetitivo.** 2022. 60 f. Dissertação (Mestrado) – Universidade Federal de Viçosa, Departamento de Biologia Animal, 2022.



SOSA-CALVO, J.; SCHULTZ, T. R. Three remarkable new fungus-growing ant species of the genus *Myrmicocrypta*, with a reassessment of the characters that define the genus and its position within the Attini. **Annals of the Entomological Society of America**, v. 103, p. 181–195, 2010.

TEIXEIRA, G. A. *et al.* Karyotype conservation and genomic organization of repetitive sequences in the leaf-cutting ant *Atta cephalotes* (Linnaeus, 1758) (Formicidae: Myrmicinae). **Genome**, v. 65, n. 10, p. 525–535, 2022. DOI: <https://doi.org/10.1139/gen-2021-0129>.

TEIXEIRA, M. L. L. *et al.* Cariótipo da formiga *Tapinoma melanocephalum* Fabricius, 1793 coletada em Oiapoque, Amapá, Amazônia. In: CONGRESSO BRASILEIRO INTERDISCIPLINAR EM CIÊNCIA E TECNOLOGIA, 2025, Diamantina. **Anais [...]** Diamantina: Even3, 2025. Disponível em: <https://www.even3.com.br/anais/congresso-brasileiro-interdisciplinar-ciencia-tecnologia/1197994>. Acesso em: jan. 2026.

TROYA, A.; LATTKE, J. *Neoponera* Emery, 1901 (Hymenoptera: Formicidae) revisited: 1. The *N. laevigata* species-group. **Insect Systematics & Evolution**, v. 54, n. 1, p. 1–76, 2022.

VÄÄNÄNEN, S.; VEPSÄLÄINEN, K.; VEPSÄLÄINEN, V. *Technomyrmex vitiensis* Mann, 1921 (Hymenoptera, Formicidae, Dolichoderinae), a new exotic tramp ant in Finland. **Sahlbergia**, v. 24, n. 1, p. 14–19, 2018.

VASCONCELOS, H. L. *et al.* Patterns of ant species diversity and turnover across 2000 km of Amazonian floodplain forest. **Journal of Biogeography**, v. 37, p. 432–440, 2010. DOI: <https://doi.org/10.1111/j.1365-2699.2009.02230.x>.

VICENTE, R. E.; PRADO, L. P.; IZZO, T. J. Amazon rainforest ant-fauna of Parque Estadual do Cristalino: understory and ground-dwelling ants. **Sociobiology**, v. 63, n. 3, p. 894–908, 2016. DOI: <https://doi.org/10.13102/sociobiology.v63i3.1043>.

VIEIRA, G. de D. *et al.* Bactérias Gram positivas veiculadas por formigas em ambiente hospitalar de Porto Velho, Estado de Rondônia, Brasil. **Revista Pan-Amazônica de Saúde**, v. 4, n. 3, p. 33–36, 2013. DOI: <https://dx.doi.org/10.5123/S2176-62232013000300005>.

VINHA, G. G. **Sistemática molecular de *Atta laevigata* (Smith, 1858) e *Acromyrmex balzani* (Emery, 1890)**. 2007. 133 f. Dissertação (Mestrado) – Universidade Estadual Paulista, Instituto de Biociências, Rio Claro, 2007.

WARD, P. S. A review of the *Pseudomyrmex ferrugineus* and *Pseudomyrmex goeldii* species groups: acacia-ants and relatives (Hymenoptera: Formicidae). **Zootaxa**, v. 4227, n. 4, p. 524–542, 2017.

WARD, P. S. Ants. **Current Biology**, v. 16, p. R152–R155, 2006.

WARD, P. S. Systematics, biogeography and host plant associations of the *Pseudomyrmex viduus* group (Hymenoptera: Formicidae), *Triplaris* and *Tachigali* inhabiting ants. **Zoological Journal of the Linnean Society**, v. 126, p. 451–540, 1999.

WATKINS, J. F. II. **The identification and distribution of new world army ants (Dorylinae: Formicidae)**. Waco, Texas: Baylor University Press, 1976. 102 p.